arXiv:hep-th/0103213v1 26 Mar 2001

D-branes and twisted K-theory

Yuri Malyuta

Institute for Nuclear Research
National Academy of Sciences of Ukraine
03022 Kiev, Ukraine
e-mail: interdep@kinr.kiev.ua

Abstract

Topological charges of the $D6$-brane in the presence of a Neveu-Schwarz B-field are computed by methods of twisted K-theory.

Keywords: D-branes, Neveu-Schwarz B-field, Twisted K-theory,
Extensions of C^*-algebras.

1 Introduction

D-branes are topological solitons, which charges are described by Grothendieck K-groups [1]. We begin our consideration with instantons [2] as they ideologically remind D-branes.

The self-dual equation for the $SU(2)$ instanton has the form

$$F_{\mu\nu} = F^{*}_{\mu\nu} \ , \tag{1.1}$$

where Yang-Mills strengths and potentials are defined as follows

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu], \quad F^{*}_{\mu\nu} = \frac{1}{2}\varepsilon_{\mu\nu\lambda\kappa}F_{\lambda\kappa}, \quad \varepsilon_{0123} = 1,$$

$$F_{\mu\nu} = \frac{1}{2i}\sigma^a F^a_{\mu\nu}(x), \quad A_\mu = \frac{1}{2i}\sigma^a A^a_\mu(x),$$

$\sigma^a/2$ are generators of the $SU(2)$ group, $x = (x_0, x_1, x_2, x_3)$ are coordinates in Euclidean space.

't Hooft [3] has found the $5k$ parameter solution of the equation (1.1)

$$A_\mu = \frac{1}{2i}\eta_{\mu\nu}\partial_\nu ln\phi \ ,$$

where $$\eta_{\mu\nu} = (\varepsilon_{oa\mu\nu} + \delta_{a\mu}\delta_{o\nu} - \delta_{a\nu}\delta_{o\mu})\sigma^a \ ,$$

$$\phi = 1 + \sum_{i=1}^{k}\frac{\lambda_i^2}{(x_\nu - y_{\nu i})^2} \ , \quad k = \frac{1}{16\pi^2}\int d^4x Tr(F_{\mu\nu}F^{*}_{\mu\nu}) \ ,$$

λ_i and $y_{\nu i}$ are parameters, k is the topological charge.

It was shown in the works of Atiyah, Hitchin, Drinfeld and Manin [4, 5] that dynamics of the equation (1.1) is encoded in the principal bundle

$$\begin{tikzcd} SU(2) \arrow[r] & P \arrow[d] \\ & S^4 \end{tikzcd} \tag{1.2}$$

and in the associated vector bundle

$$\begin{tikzcd} \mathbb{C}^2 \arrow[r] & P \arrow[d] \\ & S^4 \end{tikzcd} \tag{1.3}$$

The potential A_μ is a connection of the vector bundle (1.3), therefore it is possible to calculate it by methods of algebraic geometry. For this purpose following the techniques of Horrocks and Barth [6], it is necessary to calculate the orthonormalized basis of sections Y^p of the vector bundle (1.3). Then the potential A_μ will be equal to

$$A_\mu^{pq} = (Y^p, \partial_\mu Y^q) \ .$$

The topological charge k is an element of the homotopy group

$$\pi_3(SU(2)) = \mathbb{Z} \ ,$$

characterizing the principal bundle (1.2).

In the next sections we shall apply the algebro-geometric approach to the description of D-branes.

2 Twisted K-theory

Let us consider the following principal bundles describing D-branes [7]

$$\begin{tikzcd}
U(n) \arrow[r] & P_H \arrow[d] \\
 & X
\end{tikzcd} \tag{2.1}$$

$$\begin{tikzcd}
SU(n)/\mathbb{Z}_n \arrow[r] & P_H \arrow[d] \\
 & X
\end{tikzcd} \tag{2.2}$$

$$\begin{tikzcd}
\lim\limits_{n\to\infty} SU(n)/\mathbb{Z}_n \arrow[r] & P_H \arrow[d] \\
 & X
\end{tikzcd} \tag{2.3}$$

where X is a compact manifold.

This bundles are characterized by the Dixmier-Douady invariant [8]

$$[H] \in H^3(X,\mathbb{Z}) \ ,$$

which represents the strength of a Neveu-Schwarz B-field [9] interacting with D-branes

$$H_{\mu\nu\rho} = \partial_\mu B_{\nu\rho} + \partial_\nu B_{\rho\mu} + \partial_\rho B_{\mu\nu} \ .$$

For the bundle (2.1)

$$[H] = 0 \ , \quad \text{id est} \quad H_{\mu\nu\lambda} = 0 \ , \quad B_{\mu\nu} = 0 \ ;$$

for the bundle (2.2)

$$n[H] = 0 \ , \quad \text{id est} \quad H_{\mu\nu\lambda} = 0 \ , \quad B_{\mu\nu} \neq 0 \ ;$$

for the bundle (2.3)

$$[H] \neq 0 \ , \quad \text{id est} \quad H_{\mu\nu\lambda} \neq 0 \ , \quad B_{\mu\nu} \neq 0 \ .$$

The vector bundles associated with the principal bundles (2.1), (2.2), (2.3) are given by [7]

$$E_H = P_H \times \mathbb{C}^n \ , \quad \text{where} \quad Aut(\mathbb{C}^n) = U(n) \ ; \tag{2.4}$$
$$E_H = P_H \times M_n(\mathbb{C}) \ , \quad \text{where} \quad Aut(M_n(\mathbb{C})) = SU(n)/\mathbb{Z}_n \ ; \tag{2.5}$$
$$E_H = P_H \times \mathcal{K} \ , \quad \text{where} \quad Aut(\mathcal{K}) = \lim_{n\to\infty} SU(n)/\mathbb{Z}_n \ ; \tag{2.6}$$

$M_n(\mathbb{C})$ is the algebra of $n \times n$ matrices, $\mathcal{K}$ is the algebra of compact operators.

The space of sections of the vector bundle (2.4) is the algebra of continuous functions $C_0(X)$;

the space of sections of the vector bundle (2.5) is the C^*-algebra $C_0(X, E_H)$, which is Morita-equivalent to the Azumaya algebra [10];

the space of sections of the vector bundle (2.6) is the Rosenberg C^*-algebra $C_0(X, E_H)$ [11].

Let us consider the short exact sequences [12]

$$0 \to \mathcal{K} \to \mathcal{B} \to C_0(X, E_H) \to 0 \tag{2.7}$$

$$0 \to \mathcal{K} \to \mathcal{B}' \to C_0(X, E_H) \to 0 \tag{2.8}$$

determining extensions of the algebra $C_0(X, E_H)$ by the algebra $\mathcal{K}$.

If in the commutative diagrams

$$\begin{tikzcd}
0 \arrow[r] & \mathcal{K} \arrow[r] \arrow[d] & \mathcal{B} \arrow[r] \arrow[d] & C_0(X, E_H) \arrow[r] \arrow[d, "\tau_1"] & 0 \\
0 \arrow[r] & \mathcal{K} \arrow[r] & B(\mathcal{H}) \arrow[r, "\pi"] & \mathcal{Q}(\mathcal{H}) \arrow[r] & 0
\end{tikzcd}$$

$$\begin{tikzcd}
0 \arrow[r] & \mathcal{K} \arrow[r] \arrow[d] & \mathcal{B}' \arrow[r] \arrow[d] & C_0(X, E_H) \arrow[r] \arrow[d, "\tau_2"] & 0 \\
0 \arrow[r] & \mathcal{K} \arrow[r] & B(\mathcal{H}) \arrow[r, "\pi"] & \mathcal{Q}(\mathcal{H}) \arrow[r] & 0
\end{tikzcd}$$

(where $B(\mathcal{H})$ is the algebra of bounded operators on a Hilbert space $\mathcal{H}$, $\mathcal{Q}(\mathcal{H})$ denotes the Calkin algebra) morphisms are related by

$$\tau_2(s) = \pi(U)\tau_1(s)\pi(U)^*$$

(s $\in C_0(X, E_H)$, U is a unitary operator on $\mathcal{H}$), then two extensions (2.7) and (2.8) are unitarily equivalent.

Let $Ext(X, [H])$ denote the set of unitary equivalence classes of extensions of $C_0(X, E_H)$ by $\mathcal{K}$ modulo splitting extensions. Then twisted K-groups are defined as follows [12]

$$K_j(X, [H]) = Ext(X \times \mathbb{R}^j,\ p_1^*[H]), \quad j \in \mathbb{N}\ ,$$

where $p_1 : X \times \mathbb{R}^j \to X$ denotes projection onto the first factor. The Bott periodicity theorem asserts:

$$K_j(X, [H]) = \begin{cases} K_0(X, [H]) & \text{for even } j\ , \\ K_1(X, [H]) & \text{for odd } j\ . \end{cases}$$

3 Computation of twisted K-groups

A short exact sequence of C^*-algebras

$$0 \to \mathcal{A} \to \mathcal{B} \to C_0(X, E_H) \to 0$$

induces a hexagon exact sequence of twisted K-groups [12]

$$\begin{tikzcd} K_0(X, [H]) \arrow[r] & K_0(\mathcal{B}) \arrow[r] & K_0(\mathcal{A}) \arrow[d] \\ K_1(\mathcal{A}) \arrow[u] & K_1(\mathcal{B}) \arrow[l] & K_1(X, [H]) \arrow[l] \end{tikzcd} \tag{3.1}$$

The hexagon sequence (3.1) enables us to compute twisted K-groups.

Let $X = S^3$. We will compute twisted groups $K_0(S^3, n[H])$ and $K_1(S^3, n[H])$ for the case $H_{\mu\nu\lambda} = 0$, $B_{\mu\nu} \neq 0$. There is the short exact sequence

$$0 \to C_0(S^2) \otimes \mathcal{K} \to C_0(U_1) \otimes \mathcal{K} \ \oplus \ C_0(U_2) \otimes \mathcal{K} \to$$

$$\to C_0(S^3, E_{n[H]}) \to 0$$

(where $\{U_1, U_2\}$ is the open cover of S^3 by hemispheres) , which induces the hexagon exact sequence of K-groups

$$\begin{tikzcd}
K_0(S^3, n[H]) \arrow[r] & K_0(U_1) \oplus K_0(U_2) \arrow[r] & K_0(S^2) \arrow[d] \\
K_1(S^2) \arrow[u] & K_1(U_1) \oplus K_1(U_2) \arrow[l] & K_1(S^3, n[H]) \arrow[l]
\end{tikzcd} \tag{3.2}$$

Since

$$K_0(S^2) = \mathbb{Z} \ , \qquad K_0(U_1) \oplus K_0(U_2) = \mathbb{Z} \ ,$$

$$K_1(S^2) = 0 \ , \qquad K_1(U_1) \oplus K_1(U_2) = 0 \ ,$$

the sequence (3.2) reduces to the exact sequence

$$0 \to K_0(S^3, n[H]) \to \mathbb{Z} \to \mathbb{Z} \to K_1(S^3, n[H]) \to 0 \tag{3.3}$$

Reduction of the sequence (3.3) to the exact sequence

$$0 \to \mathbb{Z} \to \mathbb{Z} \to \mathbb{Z}_n \to 0$$

yields the following result:

$$K_0(S^3, n[H]) = 0 \ , \qquad K_1(S^3, n[H]) = \mathbb{Z}_n \ .$$

The group $K_1(S^3, n[H]) = \mathbb{Z}_n$ determines topological charges of the $D6$-brane for the case $H_{\mu\nu\lambda} = 0, \quad B_{\mu\nu} \neq 0$.
For comparision we present the Dp-brane spectrum [1] for the case $H_{\mu\nu\lambda} = 0, \quad B_{\mu\nu} = 0$:

Dp	$D9$	$D8$	$D7$	$D6$	$D5$	$D4$	$D3$	$D2$	$D1$	$D0$	$D(-1)$
S^{9-p}	S^0	S^1	S^2	S^3	S^4	S^5	S^6	S^7	S^8	S^9	S^{10}
$K_0(S^{9-p})$	$\mathbb{Z}$	0	$\mathbb{Z}$	0	$\mathbb{Z}$	0	$\mathbb{Z}$	0	$\mathbb{Z}$	0	$\mathbb{Z}$
$K_1(S^{9-p})$	0	$\mathbb{Z}$	0	$\mathbb{Z}$	0	$\mathbb{Z}$	0	$\mathbb{Z}$	0	$\mathbb{Z}$	0

Acknowledgement

This material was presented in the lecture given by the author at the Institute of Mathematics (Kiev, Ukraine). The author thanks audience for questions and comments.

References

[1] E. Witten, *D-branes and K-theory*, hep-th/9810188.

[2] E. Corrigan, *Self-dual solutions to Euclidean Yang-Mills equations*, Phys. Reports **49** (1979) 95.

[3] R. Jackiw, C. Nohl and C. Rebbi, *Conformal properties of pseudoparticle configurations*, Phys. Rev. **D15** (1977) 1642.

[4] V.G. Drinfeld and Yu.I. Manin, *Self-dual Yang-Mills fields on the sphere*, Funkt. Anal. Priloz. **12**, N2 (1978) 78.

[5] M.F. Atiyah, N.J. Hitchin, V.G. Drinfeld and Yu.I. Manin, *Construction of instantons*, Phys. Lett. **65A** (1978) 185.

[6] W. Barth, *Moduli of vector bundles on the projective plane*, Inventiones math. **42** (1977) 63.

[7] P. Bouwknegt and V. Mathai, *D-branes, B-fields and twisted K-theory*, hep-th/0002023.

[8] J. Dixmier and A. Douady, *Champs continues d'espaces hilbertiens at de C^*-algebres*, Bull. Soc. Math. France **91** (1963) 227.

[9] E. Witten, *Overview of K-theory applied to strings*, hep-th/0007175.

[10] A. Kapustin, *D-branes in a topologically nontrivial B-field*, hep-th/9909089.

[11] J. Rosenberg, *Continuous trace algebras from the bundle theoretic point of view*, Jour. Austr. Math. Soc. **47** (1989) 368.

[12] V. Mathai and I.M. Singer, *Twisted K-homology theory, twisted Ext-theory*, hep-th/0012046.